arXiv:1201.6226v1 [math.FA] 25 Jan 2012

NEW INTEGRAL INEQUALITIES VIA (α, m)-CONVEXITY AND QUASI-CONVEXITY

WENJUN LIU

Abstract. In this paper, we establish some new integral inequalities for $(\alpha, m)-$convex functions and quasi-convex functions, respectively. Our results in special cases recapture known results.

1. INTRODUCTION

Let I be on interval in R. Then $f : I \to R$ is said to be convex (see [17, P.1]) if

$$f\left(tx + (1-t)\, y\right) \le tf\left(x\right) + (1-t)\, f\left(y\right)$$

holds for all $x, y \in I$ and $t \in [0,1]$.

In [27], Toader defined m-convexity as follows:

Definition 1. *The function $f : [0,b] \to R$, $b > 0$ is said to be m-convex, where $m \in [0,1]$, if*

$$f\left(tx + m\left(1-t\right) y\right) \le tf\left(x\right) + m\left(1-t\right) f\left(y\right)$$

holds for all $x, y \in [0,b]$ and $t \in [0,1]$. We say that f is $m-$concave if $-f$ is $m-$convex.

In [18], Miheşan defined $(\alpha, m)-$ convexity as follows:

Definition 2. *The function $f : [0,b] \to \mathbb{R}$, $b > 0$, is said to be $(\alpha, m)-$ convex, where $(\alpha, m) \in [0,1]^2$, if*

$$f(tx + m(1-t)y) \le t^{\alpha} f(x) + m(1-t^{\alpha}) f(y)$$

holds for all $x, y \in [0,b]$ and $t \in [0,1]$.

Denote by $K_m^{\alpha}(b)$ the class of all $(\alpha, m)-$convex functions on $[0,b]$ for which $f(0) \le 0$. It can be easily seen that for $(\alpha, m) = (1, m)$, $(\alpha, m)-$ convexity reduces to $m-$ convexity and for $(\alpha, m) = (1,1)$, $(\alpha, m)-$ convexity reduces to the concept of usual convexity defined on $[0,b]$, $b > 0$. For recent results and generalizations concerning $m-$convex and $(\alpha, m)-$convex functions see [4, 6, 10, 19, 21, 26].

We recall that the notion of quasi-convex functions generalizes the notion of convex functions. More precisely, a function $f : [a,b] \to \mathbb{R}$ is said to be quasi-convex on $[a,b]$ if

$$f(\lambda x + (1-\lambda)y) \le \max\{f(x), f(y)\}$$

holds for any $x, y \in [a,b]$ and $\lambda \in [0,1]$. Clearly, any convex function is a quasi-convex function. Furthermore, there exist quasi-convex functions which are not convex (see [14]).

One of the most famous inequalities for convex functions is Hadamard's inequality. This double inequality is stated as follows: Let f be a convex function on some nonempty interval $[a,b]$ of real line $\mathbb{R}$, where $a \ne b$. Then

$$f\left(\frac{a+b}{2}\right) \le \frac{1}{b-a}\int_a^b f(x)dx \le \frac{f\left(a\right)+f\left(b\right)}{2}. \tag{1.1}$$

Hadamard's inequality for convex functions has received renewed attention in recent years and a remarkable variety of refinements and generalizations have been found (see, for example, [1]-[19], [22]-[26], [28]). In [4], Bakula et al. establish several Hadamard type inequalities for differentiable $m-$convex and $(\alpha, m)-$convex functions.

2000 *Mathematics Subject Classification.* 26D15, 26A51, 39B62.

Key words and phrases. Hermite's inequality, Hölder's inequality, (α, m)-convexity, quasi-convexity.

Recently, Ion [14] established two estimates on the Hermite-Hadamard inequality for functions whose first derivatives in absolute value are quasi-convex. Namely, he obtained the following results:

Theorem 1.1. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I, $a, b \in I$ with $a < b$. If $|f'|$ is quasi-convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(u)du\right| \le \frac{b-a}{4}\left\{\max|f'(a)|, |f'(b)|\right\}.$$

Theorem 1.2. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I, $a, b \in I$ with $a < b$ and let $p > 1$. If $|f'|^{\frac{p}{p-1}}$ is quasi-convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(u)du\right| \le \frac{b-a}{2(p+1)^{\frac{1}{p}}}\left(\max\left\{|f'(a)|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}}\right\}\right)^{\frac{p-1}{p}}.$$

In [2], Alomari et al. obtained the following result.

Theorem 1.3. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I, $a, b \in I$ with $a < b$ and let $q \ge 1$. If $|f'|^q$ is quasi-convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(u)du\right| \le \frac{b-a}{4}\left(\max\left\{|f'(a)|^q, |f'(b)|^q\right\}\right)^{\frac{1}{q}}.$$

In [20], Özdemir et al. used the following lemma in order to establish several integral inequalities via some kinds of convexity.

Lemma 1.1. *Let $f : [a, b] \subset [0, \infty) \to \mathbb{R}$ be continuous on $[a, b]$ such that $f \in L([a, b])$, $a < b$. Then the equality*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx = (b-a)^{p+q+1}\int_0^1 (1-t)^p t^q f(ta+(1-t)b)dt \tag{1.2}$$

holds for some fixed $p, q > 0$.

Especially, Özdemir et al. [20] discussed the following new results connecting with $m-$convex function and quasi-convex function, respectively:

Theorem 1.4. *Let $f : [a, b] \to \mathbb{R}$ be continuous on $[a, b]$ such that $f \in L([a, b])$, $0 \le a < b < \infty$. If f is $m-$convex on $[a, b]$, for some fixed $m \in (0, 1]$ and $p, q > 0$, then*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\min\left\{\beta(q+2,p+1)f(a)+m\beta(q+1,p+2)f\left(\frac{b}{m}\right),\right.\\ &\left.\beta(q+1,p+2)f(b)+m\beta(q+2,p+1)f\left(\frac{a}{m}\right)\right\},\end{aligned}\tag{1.3}$$

where $\beta(x, y)$ is the Euler Beta function.

Theorem 1.5. *Let $f : [a, b] \to \mathbb{R}$ be continuous on $[a, b]$ such that $f \in L([a, b])$, $0 \le a < b < \infty$. If f is quasi-convex on $[a, b]$, then for some fixed $p, q > 0$, we have*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\max\{f(a), f(b)\}\beta(p+1, q+1).\tag{1.4}$$

The aim of this paper is to establish some new integral inequalities like those given in Theorems 1.4 and 1.5 for $(\alpha, m)-$convex functions (Section 2) and quasi-convex functions (Section 3), respectively. Our results in special cases recapture Theorems 1.4 and 1.5, respectively. That is, this study is a continuation and generalization of [20].

2. New integral inequalities for $(\alpha,m)-$ convex functions

Theorem 2.1. *Let $f:[a,b]\to\mathbb{R}$ be continuous on $[a,b]$ such that $f\in L([a,b])$, $0\le a<b<\infty$. If f is $(\alpha,m)-$convex on $[a,b]$, for some fixed $(\alpha,m)\in(0,1]^2$ and $p,q>0$, then*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\min\left\{\beta(q+\alpha+1,p+1)f(a)+m[\beta(q+1,p+1)-\beta(q+\alpha+1,p+1)]f\left(\frac{b}{m}\right),\right.\\ &\left.\beta(q+1,p+\alpha+1)f(b)+m[\beta(p+1,q+1)-\beta(q+1,p+\alpha+1)]f\left(\frac{a}{m}\right)\right\}, \qquad (2.1)\end{aligned}$$

where $\beta(x,y)$ is the Euler Beta function.

Proof. Since f is $(\alpha,m)-$convex on $[a,b]$, we know that for every $t\in[0,1]$

$$f(ta+(1-t)b)=f\left(ta+m(1-t)\frac{b}{m}\right)\le t^\alpha f(a)+m\left(1-t^\alpha\right)f\left(\frac{b}{m}\right). \qquad (2.2)$$

Using Lemma 1.1, with $x=ta+(1-t)b$, then we have

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\int_0^1(1-t)^pt^q\left(t^\alpha f(a)+m\left(1-t^\alpha\right)f\left(\frac{b}{m}\right)\right)dt\\ =&(b-a)^{p+q+1}\left[f(a)\int_0^1(1-t)^pt^{q+\alpha}dt+mf\left(\frac{b}{m}\right)\int_0^1(1-t)^pt^q\left(1-t^\alpha\right)dt\right].\end{aligned}$$

Now, we will make use of the Beta function which is defined for $x,y>0$ as

$$\beta(x,y)=\int_0^1 t^{x-1}(1-t)^{y-1}dt.$$

It is known that

$$\int_0^1 t^{q+\alpha}(1-t)^pdt=\beta(q+\alpha+1,p+1),$$

$$\begin{aligned}\int_0^1(1-t)^pt^q\left(1-t^\alpha\right)dt=&\int_0^1 t^q(1-t)^pdt-\int_0^1 t^{q+\alpha}(1-t)^pdt\\ =&\beta(q+1,p+1)-\beta(q+\alpha+1,p+1)].\end{aligned}$$

Combining all obtained equalities we get

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left\{\beta(q+\alpha+1,p+1)f(a)+m[\beta(q+1,p+1)-\beta(q+\alpha+1,p+1)]f\left(\frac{b}{m}\right)\right\}. \qquad (2.3)\end{aligned}$$

If we choose $x=tb+(1-t)a$, analogously we obtain

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left\{\beta(q+1,p+\alpha+1)f(b)+m[\beta(q+1,p+1)-\beta(q+1,p+\alpha+1)]f\left(\frac{a}{m}\right)\right\}. \qquad (2.4)\end{aligned}$$

Thus, by (2.3) and (2.4) we obtain (2.1), which completes the proof. □

Remark 1. *As a special case of Theorem 2.1 for $\alpha = 1$, that is for f be $m-$convex on $[a,b]$, we recapture Theorem 1.4 due to the fact that*

$$\begin{aligned}\beta(q+1,p+1)-\beta(q+2,p+1) =&\beta(q+1,p+1)-\frac{q+1}{p+q+2}\beta(q+1,p+1)\\ =&\frac{p+1}{p+q+2}\beta(q+1,p+1)=\beta(q+1,p+2)\end{aligned}$$

and

$$\beta(q+1,p+1)-\beta(q+1,p+\alpha+1)=\beta(q+2,p+1).$$

Corollary 2.1. *In Theorem 2.1, if $p = q$, then (2.1) reduces to*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^p f(x)dx\\ \le&(b-a)^{2p+1}\min\left\{\beta(p+\alpha+1,p+1)f(a)+m[\beta(p+1,p+1)-\beta(p+\alpha+1,p+1)]f\left(\frac{b}{m}\right),\right.\\ &\left.\beta(p+1,p+\alpha+1)f(b)+m[\beta(p+1,p+1)-\beta(p+1,p+\alpha+1)]f\left(\frac{a}{m}\right)\right\}.\end{aligned}$$

Theorem 2.2. *Let $f : [a,b] \to \mathbb{R}$ be continuous on $[a,b]$ such that $f \in L([a,b])$, $0 \le a < b < \infty$ and let $k > 1$. If $|f|^{\frac{k}{k-1}}$ is $(\alpha,m)-$convex on $[a,b]$, for some fixed $(\alpha,m) \in (0,1]^2$ and $p,q > 0$, then*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&\frac{(b-a)^{p+q+1}}{(\alpha+1)^{\frac{k-1}{k}}}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}}\min\left\{\left[|f(a)|^{\frac{k}{k-1}}+\alpha m\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}},\right.\\ &\left.\left[|f(b)|^{\frac{k}{k-1}}+\alpha m\left|f\left(\frac{a}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}}\right\}.\end{aligned}\tag{2.5}$$

Proof. Since $|f|^{\frac{k}{k-1}}$ is $(\alpha,m)-$convex on $[a,b]$ we know that for every $t \in [0,1]$

$$\begin{aligned}|f(ta+(1-t)b)|^{\frac{k}{k-1}}=&\left|f\left(ta+m(1-t)\frac{b}{m}\right)\right|^{\frac{k}{k-1}}\\ \le&t^\alpha|f(a)|^{\frac{k}{k-1}}+m\left(1-t^\alpha\right)\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}.\end{aligned}$$

Using Lemma 1.1, with $x = ta + (1-t)b$, then we have

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left[\int_0^1(1-t)^{kp}t^{kq}dt\right]^{\frac{1}{k}}\left[\int_0^1|f(ta+(1-t)b)|^{\frac{k}{k-1}}dt\right]^{\frac{k-1}{k}}\\ \le&(b-a)^{p+q+1}\left[\beta(kq+1,kp+1)\right]^{\frac{1}{k}}\left[\int_0^1 t^\alpha|f(a)|^{\frac{k}{k-1}}dt+m\int_0^1\left(1-t^\alpha\right)\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}dt\right]^{\frac{k-1}{k}}\\ =&(b-a)^{p+q+1}\left[\beta(kq+1,kp+1)\right]^{\frac{1}{k}}\left[\frac{1}{\alpha+1}|f(a)|^{\frac{k}{k-1}}+m\frac{\alpha}{\alpha+1}\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}}.\end{aligned}$$

If we choose $x = tb + (1-t)a$, analogously we obtain

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}}\left[\frac{1}{\alpha+1}|f(b)|^{\frac{k}{k-1}}+m\frac{\alpha}{\alpha+1}\left|f\left(\frac{a}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}},\end{aligned}$$

which completes the proof. □

Corollary 2.2. *In Theorem 2.2, if $p = q$, then (2.5) reduces to*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^p f(x)dx\\ \le&\frac{(b-a)^{2p+1}}{(\alpha+1)^{\frac{k-1}{k}}}\left[\beta(kp+1,kp+1)\right]^{\frac{1}{k}}\min\left\{\left[|f(a)|^{\frac{k}{k-1}}+\alpha m\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}},\right.\\ &\left.\left[|f(b)|^{\frac{k}{k-1}}+\alpha m\left|f\left(\frac{a}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}}\right\}.\end{aligned}$$

Corollary 2.3. *In Theorem 2.2, if $\alpha = 1$, i.e., if $|f|^{\frac{k}{k-1}}$ is $m-$convex on $[a,b]$, then (2.5) reduces to*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&\frac{(b-a)^{p+q+1}}{2^{\frac{k-1}{k}}}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}}\min\left\{\left[|f(a)|^{\frac{k}{k-1}}+m\left|f\left(\frac{b}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}},\right.\\ &\left.\left[|f(b)|^{\frac{k}{k-1}}+m\left|f\left(\frac{a}{m}\right)\right|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}}\right\}.\end{aligned}$$

Remark 2. *As a special case of Corollary 2.3 for $m = 1$, that is for $|f|^{\frac{k}{k-1}}$ be convex on $[a,b]$, we get*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx\le\frac{(b-a)^{p+q+1}}{2^{\frac{k-1}{k}}}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}}\left[|f(a)|^{\frac{k}{k-1}}+|f(b)|^{\frac{k}{k-1}}\right]^{\frac{k-1}{k}}.$$

Theorem 2.3. *Let $f : [a,b] \to \mathbb{R}$ be continuous on $[a,b]$ such that $f \in L([a,b])$, $0 \le a < b < \infty$ and let $l \ge 1$. If $|f|^l$ is $(\alpha,m)-$convex on $[a,b]$, for some fixed $(\alpha,m) \in (0,1]^2$ and $p, q > 0$, then*

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left[\beta(p+1,q+1)\right]^{\frac{l-1}{l}}\\ &\times\min\left\{\left[\beta(q+\alpha+1,p+1)|f(a)|^l+m[\beta(q+1,p+1)-\beta(q+\alpha+1,p+1)]\left|f\left(\frac{b}{m}\right)\right|^l\right]^{\frac{1}{l}},\right.\\ &\left.\left[\beta(q+1,p+\alpha+1)|f(b)|^l+m[\beta(q+1,p+1)-\beta(q+1,p+\alpha+1)]\left|f\left(\frac{a}{m}\right)\right|^l\right]^{\frac{1}{l}}\right\}.\end{aligned}\tag{2.6}$$

Proof. Since $|f|^l$ is $(\alpha,m)-$convex on $[a,b]$, we know that for every $t \in [0,1]$

$$|f(ta+(1-t)b)|^l=\left|f\left(ta+m(1-t)\frac{b}{m}\right)\right|^l\le t^\alpha|f(a)|^l+m\left(1-t^\alpha\right)\left|f\left(\frac{b}{m}\right)\right|^l.$$

Using Lemma 1.1, with $x = ta + (1-t)b$, then we have

$$\begin{aligned}
&\int_a^b (x-a)^p (b-x)^q f(x)dx \\
=&(b-a)^{p+q+1} \int_0^1 \left[(1-t)^p t^q\right]^{\frac{l-1}{l}} \left[(1-t)^p t^q\right]^{\frac{1}{l}} f(ta+(1-t)b)dt \\
\leq&(b-a)^{p+q+1} \left[\int_0^1 (1-t)^p t^q dt\right]^{\frac{l-1}{l}} \left[\int_0^1 (1-t)^p t^q |f(ta+(1-t)b)|^l dt\right]^{\frac{1}{l}} \\
\leq&(b-a)^{p+q+1} \left[\beta(q+1,p+1)\right]^{\frac{l-1}{l}} \\
&\times \left[\beta(q+\alpha+1,p+1)|f(a)|^l + m[\beta(q+1,p+1) - \beta(q+\alpha+1,p+1)] \left|f\left(\frac{b}{m}\right)\right|^l\right]^{\frac{1}{l}}.
\end{aligned}$$

If we choose $x = tb + (1-t)a$, analogously we obtain

$$\begin{aligned}
&\int_a^b (x-a)^p (b-x)^q f(x)dx \\
\leq&(b-a)^{p+q+1} \left[\beta(p+1,q+1)\right]^{\frac{l-1}{l}} \\
&\times \left[\beta(q+1,p+\alpha+1)|f(b)|^l + m[\beta(q+1,p+1) - \beta(q+1,p+\alpha+1)] \left|f\left(\frac{a}{m}\right)\right|^l\right]^{\frac{1}{l}},
\end{aligned}$$

which completes the proof. □

Corollary 2.4. *In Theorem 2.3, if $p = q$, then* (2.6) *reduces to*

$$\begin{aligned}
&\int_a^b (x-a)^p (b-x)^p f(x)dx \\
\leq&(b-a)^{2p+1} \left[\beta(p+1,p+1)\right]^{\frac{l-1}{l}} \\
&\times \min\left\{\left[\beta(p+\alpha+1,p+1)|f(a)|^l + m[\beta(p+1,p+1) - \beta(p+\alpha+1,p+1)] \left|f\left(\frac{b}{m}\right)\right|^l\right]^{\frac{1}{l}},\right. \\
&\left.\left[\beta(p+1,p+\alpha+1)|f(b)|^l + m[\beta(p+1,p+1) - \beta(p+1,p+\alpha+1)] \left|f\left(\frac{a}{m}\right)\right|^l\right]^{\frac{1}{l}}\right\}.
\end{aligned}$$

Corollary 2.5. *In Theorem 2.3, if $\alpha = 1$, i.e., if $|f|^l$ is $m-$convex on $[a,b]$, then* (2.6) *reduces to*

$$\begin{aligned}
&\int_a^b (x-a)^p (b-x)^q f(x)dx \\
\leq&(b-a)^{p+q+1} \left[\beta(p+1,q+1)\right]^{\frac{l-1}{l}} \min\left\{\left[\beta(q+2,p+1)|f(a)|^l + m\beta(q+1,p+2) \left|f\left(\frac{b}{m}\right)\right|^l\right]^{\frac{1}{l}},\right. \\
&\left.\left[\beta(q+1,p+2)|f(b)|^l + m\beta(q+2,p+1) \left|f\left(\frac{a}{m}\right)\right|^l\right]^{\frac{1}{l}}\right\}.
\end{aligned}$$

Remark 3. *As a special case of Corollary 2.5 for $m = 1$, that is for $|f|^l$ be convex on $[a,b]$, we get*

$$\begin{aligned}
&\int_a^b (x-a)^p (b-x)^q f(x)dx \\
\leq&(b-a)^{p+q+1} \left[\beta(p+1,q+1)\right]^{\frac{l-1}{l}} \left[\beta(q+2,p+1)|f(a)|^l + \beta(q+1,p+2)\left|f(b)\right|^l\right]^{\frac{1}{l}}.
\end{aligned}$$

3. New integral inequalities for quasi-convex functions

Theorem 3.1. *Let $f:[a,b]\to\mathbb{R}$ be continuous on $[a,b]$ such that $f\in L([a,b])$, $0\le a<b<\infty$ and let $k>1$. If $|f|^{\frac{k}{k-1}}$ is quasi-convex on $[a,b]$, for some fixed $p,q>0$, then*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}}\left(\max\left\{|f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}}\right\}\right)^{\frac{k-1}{k}}. \tag{3.1}$$

Proof. By Lemma 1.1, Hölder's inequality, the definition of Beta function and the fact that $|f|^{\frac{k}{k-1}}$ is quasi-convex on $[a,b]$, we have

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ \le&(b-a)^{p+q+1}\left[\int_0^1 (1-t)^{kp}t^{kq}dt\right]^{\frac{1}{k}}\left[\int_0^1 |f(ta+(1-t)b)|^{\frac{k}{k-1}}dt\right]^{\frac{k-1}{k}}\\ \le&(b-a)^{p+q+1}\left[\beta(kq+1,kp+1)\right]^{\frac{1}{k}}\left[\int_0^1 \max\left\{|f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}}\right\}dt\right]^{\frac{k-1}{k}}\\ =&(b-a)^{p+q+1}\left[\beta(kq+1,kp+1)\right]^{\frac{1}{k}}\left[\max\left\{|f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}}\right\}\right]^{\frac{k-1}{k}},\end{aligned}$$

which completes the proof. □

Corollary 3.1. *Let f be as in Theorem 3.1. Additionally, if*

(1) *f is increasing, then we have*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}} f(b).$$

(2) *f is decreasing, then we have*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\left[\beta(kp+1,kq+1)\right]^{\frac{1}{k}} f(a).$$

Theorem 3.2. *Let $f:[a,b]\to\mathbb{R}$ be continuous on $[a,b]$ such that $f\in L([a,b])$, $0\le a<b<\infty$ and let $l\ge 1$. If $|f|^l$ is quasi-convex on $[a,b]$, for some fixed $p,q>0$, then*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\beta(p+1,q+1)\left(\max\left\{|f(a)|^l, |f(b)|^l\right\}\right)^{\frac{1}{l}}, \tag{3.2}$$

where $\beta(x,y)$ is the Euler Beta function.

Proof. By Lemma 1.1, Hölder's inequality, the definition of Beta function and the fact that $|f|^l$ is quasi-convex on $[a,b]$, we have

$$\begin{aligned}&\int_a^b (x-a)^p(b-x)^q f(x)dx\\ =&(b-a)^{p+q+1}\int_0^1 \left[(1-t)^pt^q\right]^{\frac{l-1}{l}}\left[(1-t)^pt^q\right]^{\frac{1}{l}} f(ta+(1-t)b)dt\\ \le&(b-a)^{p+q+1}\left[\int_0^1 (1-t)^pt^qdt\right]^{\frac{l-1}{l}}\left[\int_0^1 (1-t)^pt^q|f(ta+(1-t)b)|^ldt\right]^{\frac{1}{l}}\\ \le&(b-a)^{p+q+1}\left[\beta(q+1,p+1)\right]^{\frac{l-1}{l}}\left[\max\left\{|f(a)|^l, |f(b)|^l\right\}\beta(q+1,p+1)\right]^{\frac{1}{l}}\\ =&(b-a)^{p+q+1}\beta(p+1,q+1)\left(\max\left\{|f(a)|^l, |f(b)|^l\right\}\right)^{\frac{1}{l}},\end{aligned}$$

which completes the proof. □

Corollary 3.2. *Let f be as in Theorem 3.2. Additionally, if*

(1) *f is increasing, then we have*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\beta(p+1,q+1)f(b).$$

(2) *f is decreasing, then we have*

$$\int_a^b (x-a)^p(b-x)^q f(x)dx \le (b-a)^{p+q+1}\beta(p+1,q+1)f(a).$$

References

[1] M. Alomari and M. Darus, On the Hadamard's inequality for log-convex functions on the coordinates, J. Inequal. Appl. **2009**, Art. ID 283147 13 pp.

[2] M. Alomari, M. Darus and S.S. Dragomir, Inequalities of Hermite-Hadamard's type for functions whose derivatives absolute values are quasi-convex, RGMIA Res. Rep. Coll., 12 (2009), Supp., No. 14.

[3] A. G. Azpeitia, Convex functions and the Hadamard inequality, Rev. Colombiana Mat. **28** (1994), no. 1, 7–12.

[4] M. K. Bakula, M. E. Özdemir and J. Pečarić, Hadamard type inequalities for m-convex and (α,m)-convex functions, JIPAM. J. Inequal. Pure Appl. Math. **9** (2008), no. 4, Article 96, 12 pp. (electronic).

[5] M. K. Bakula and J. Pečarić, Note on some Hadamard-type inequalities, JIPAM. J. Inequal. Pure Appl. Math. **5** (2004), no. 3, Article 74, 9 pp. (electronic).

[6] M. K. Bakula, J. Pečarić and M. Ribičić, Companion Inequalities to Jensen's Inequality for m-convex and (α,m)-convex Functions, JIPAM. J. Inequal. Pure Appl. Math. **7** (2006), no. 5, Article 194, 15 pp. (electronic).

[7] C. Dinu, Hermite-Hadamard inequality on time scales, J. Inequal. Appl. **2008**, Art. ID 287947, 24 pp.

[8] S.S. Dragomir and C.E.M. Pearce, Selected Topics on Hermite-Hadamard Inequalities and Applications, *RGMIA Monographs, Victoria University,* 2000.

[9] S. S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, Appl. Math. Lett. **11** (1998), no. 5, 91–95.

[10] S. S. Dragomir, On some new inequalities of Hermite-Hadamard type for m-convex functions, Tamkang J. Math. **33** (2002), no. 1, 55–65.

[11] S. S. Dragomir and S. Fitzpatrick, The Hadamard inequalities for s-convex functions in the second sense, Demonstratio Math. **32** (1999), no. 4, 687–696.

[12] P. M. Gill, C. E. M. Pearce and J. Pečarić, Hadamard's inequality for r-convex functions, J. Math. Anal. Appl. **215** (1997), no. 2, 461–470.

[13] V. N. Huy and N. T. Chung, Some generalizations of the Fejér and Hermite-Hadamard inequalities in Hölder spaces, J. Appl. Math. Inform. **29** (2011), no. 3-4, 859–868.

[14] D. A. Ion, Some estimates on the Hermite-Hadamard inequality through quasi-convex functions, An. Univ. Craiova Ser. Mat. Inform. **34** (2007), 83–88.

[15] U. S. Kirmaci et al., Hadamard-type inequalities for s-convex functions, Appl. Math. Comput. **193** (2007), no. 1, 26–35.

[16] Z. Liu, Generalization and improvement of some Hadamard type inequalities for Lipschitzian mappings, J. Pure Appl. Math. Adv. Appl. **1** (2009), no. 2, 175–181.

[17] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and new inequalities in analysis*, Mathematics and its Applications (East European Series), 61, Kluwer Acad. Publ., Dordrecht, 1993.

[18] V. G. Miheşan, A generalization of the convexity, *Seminar on Functional Equations, Approx. and Convex.,* Cluj-Napoca (Romania) (1993)

[19] M. E. Özdemir, M. Avcı and E. Set, On some inequalities of Hermite-Hadamard type via m-convexity, Appl. Math. Lett. **23** (2010), no. 9, 1065–1070.

[20] M. E. Özdemir, E. Set and M. Alomari, Integral inequalities via several kinds of convexity, Creat. Math. Inform. **20** (2011), no. 1, 62–73.

[21] M. E. Özdemir, E. Set and M. Z. Sarıkaya, Some new Hadamard type inequalities for co-ordinated m-convex and (α,m)-convex functions, Hacet. J. Math. Stat. **40** (2011), no. 2, 219–229.

[22] J. E. Pečarić, F. Proschan and Y. L. Tong, *Convex functions, partial orderings, and statistical applications*, Mathematics in Science and Engineering, 187, Academic Press, Boston, MA, 1992.

[23] M. Z. Sarikaya, E. Set and M. E. Özdemir, On some new inequalities of Hadamard type involving h-convex functions, Acta Math. Univ. Comenian. (N.S.) **79** (2010), no. 2, 265–272.

[24] E. Set, M. E. Özdemir and S. S. Dragomir, On the Hermite-Hadamard inequality and other integral inequalities involving two functions, J. Inequal. Appl. **2010**, Art. ID 148102, 9 pp.

[25] E. Set, M. E. Özdemir and S. S. Dragomir, On Hadamard-type inequalities involving several kinds of convexity, J. Inequal. Appl. **2010**, Art. ID 286845, 12 pp.

[26] E. Set, M. Sardari, M. E. Özdemir and J. Rooin, On generalizations of the Hadamard inequality for (α,m)-convex functions, RGMIA Res. Rep. Coll., 12 (4) (2009), No. 4.

[27] G. Toader, Some generalizations of the convexity, in *Proceedings of the colloquium on approximation and optimization (Cluj-Napoca, 1985)*, 329–338, Univ. Cluj-Napoca, Cluj.
[28] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, New Hermite-Hadamard-type inequalities for convex functions (II), Comput. Math. Appl. **62** (2011), no. 1, 401–418.

(W. J. Liu) College of Mathematics and Statistics, Nanjing University of Information Science and Technology, Nanjing 210044, China
E-mail address: `wjliu@nuist.edu.cn`